
**1992 ONTARIO OPEN FIRST AID COMPETITION
TEAM SCENARIO (15 MINUTES)**

YOUR FIRST AID TEAM IS IN A STADIUM.

**THEY HAVE BEEN ASKED TO PARTICIPATE
WITH OLYMPIC TEAM MEDICAL CREWS IN
THIS YEAR'S OLYMPICS.**

**DURING THE PARADE OF COMPETITORS AN
EARTHQUAKE ROCKS THE STADIUM.**

**THE DATE, TIME AND TEMPERATURE ARE AS
YOU FIND THEM IN THIS ROOM.**

GOOD LUCK!

**ONTARIO OPEN FIRST AID COMPETITION - 1992
TEAM SCENARIO**

TOTAL SCORE: _____ / 800

(place team label here)

TIME START: _____ FINISH: _____ ELAPSED: _____

JUDGE'S NAMES (please print): _____

THE SCENARIO (15 minutes)

A first aid team has been asked to participate with olympic team medical crews in this year's olympics. During the parade of teams an earthquake rocks the stadium injuring many spectators. There are no immediate dangers in the stadium. The date, time and temperature are as inside.

THE INJURIES (Male patient only; patient is conscious and alert)

- | | |
|--|---|
| 1. Closed fractured pelvis (rt.hip, no simulation) | 5. Previous history of stroke, rt. sided weakness |
| 2. Open fracture of rt. mid-clavicle (minor bleed) | 6. Skin is pale and sweaty |
| 3. Closed fracture of rt. wrist (thumb side) | 7. Contusion on rt.shoulder |
| 4. 2" oblique laceration of lt. palm (hand) | |

JUDGES

Please assist team and patient with prompting. Patient has previous history of stroke and has rt. sided weakness. He is wearing a medic alert bracelet for same. Check the patient's position and simulation. Immediate action must be completed in part before secondary action, to receive any marks for immediate action. Immediate and secondary actions do not have to be in sequence.

(-)Judge's prompts I-Information requested A-Actions taken S-Statements made

PT. POSITION



Immediate Actions

1. Was inquiry made of hazards before further action? (there are no hazards, stadium is intact)	AS	20	0
2. Did first aider(s) call for help? (none readily available)	A	20	0
3. Did first aider(s) identify themselves? (identification as a first aider, or someone here to help is sufficient)	A	20	0
4. Was patient warned not to move?	A	20	0
5. Did first aider(s) assess for adequate airway and breathing? (airway and breathing are adequate)	A	20	0
6. Did first aider(s) assess for and control severe bleeding from palm of lt. hand?	A	20	0

Secondary Actions

7. Was a pulse (carotid or radial) taken? (min. of 15 sec. - corrected to 100/min. and weak)	A	10	0
8. Were respirations taken? (minimum of 15 sec. - corrected to 20/min. and shallow)	A	10	0
9. Was patient asked where they hurt? (pt. c/o pain down rt. side)	I	10	0

10. Was head examined for injuries?	A	10	0
11. Were both ears examined? (no blood or fluid)	A	10	0
12. Were both eyes examined? (pupils equal & reactive to light)	A	10	0
13. Was nose examined? (no blood or fluid)	A	10	0
14. Was mouth examined? (no dentures, foreign objects, bleeding, or odour)	A	10	0
15. Was jaw examined? (no difficulty moving)	A	10	0
16. Was neck examined? (no pain)	A	10	0
17. Was back (spine) examined? (no pain)	A	10	0
18. Was chest examined? (pt. c/o pain in rt. shoulder area, air entry equal and bilateral, chest sounds clear, if asked bleeding from collarbone fracture is minor)	A	10	0
19. Was abdomen examined? (no pain or rigidity)	A	10	0
20. Was pelvis examined? (pt. c/o pain rt. hip, no wetness)	A	10	0
21. Were lower extremities examined? (rt. sided weakness, no swelling, capillary refill sluggish)	A	10	0
22. Were upper extremities examined? (pt. c/o pain in rt. arm, rt. sided weakness, no swelling, capillary refill in both hands sluggish, if asked - bleeding from lt. hand is severe)	A	10	0
23. Was patient questioned re: medical history? (history of stroke resulting in rt. sided weakness for past 6 months)	I	10	0
24. Was medic alert bracelet identified?	A	10	0
25. Was patient questioned re: allergies? (none)	I	10	0
26. Was patient questioned re: medications? (takes blood thinners [Coumadin] daily)	I	10	0
27. Was patient kept in position found during examination?	A	20	0
28. Was patient SAFELY positioned to facilitate first aid for injuries? (SAFELY is defined as: patient being positioned by team, following patient assessment, using a method that provides as much support and safety as possible - if yes, then 20 marks)	A	20	0
29. Was laceration to lt. palm effectively dressed?	A	20	0
30. Was rt. mid-clavicle fracture suspected?	AS	20	0
31. Was bleeding at clavicle fracture site controlled?	A	20	0

32. Was contusion on rt. shoulder identified? (team may suspect shoulder contusion to be fracture, sprain or dislocation - that's OK)	AS	10	0
33. Was a cold compress applied to shoulder contusion?	A	10	0
34. Was rt. wrist fracture suspected?	AS	20	0
35. Was rt. arm (including wrist) effectively immobilized?	A	20	0
36. Was rt. arm (including wrist) effectively supported?	A	20	0
37. Was a pelvic (hip) fracture suspected?	AS	20	0
38. Was the pelvis effectively supported?	A	20	0
39. Was the potential for shock recognized?	AS	20	0
40. Was the patient covered to maintain body heat? (over)	A	20	0
41. Was the pulse reassessed? (minimum of 15 sec. - corrected to 100/min. and weak)	A	10	0
42. Were respirations reassessed? (minimum of 15 sec. - corrected to 20/min. and shallow)	A	10	0
43. Was the patient reassured? (at least once)	A	20	0
44. Was ambulance (EMS) called for? (will respond A.S.A.P.)	AS	20	0
<u>Recording</u>			
45. Was the patient's name recorded? (if asked - the patient's name is their real name)	A	10	0
46. Was the date and the time of the incident recorded? (if asked - the date and time are the current date and time)	A	10	0
47. Was the pulse rate recorded?	A	10	0
48. Was the respiration rate recorded?	A	10	0
49. Was the level of consciousness recorded?	A	10	0
50. Was stroke history recorded?	A	10	0
51. Was medication information recorded?	A	10	0
52. Was rt. sided weakness recorded?	A	10	0
53. Was lt. hand laceration recorded?	A	10	0
54. Was fractured rt. clavicle recorded?	A	10	0
55. Was rt. shoulder injury recorded?	A	10	0
56. Was rt. wrist fracture recorded?	A	10	0
57. Was fractured pelvis (hip) recorded?	A	10	0

58. Was lt. hand treatment recorded?

59. Was injured rt. clavicle/shoulder/wrist treatment recorded?

60. Was fractured pelvis (hip) treatment recorded?

A	10	0
A	10	0
A	10	0

Total: / 800